NJ Occupational Health Indicators



>> Hospitalizations from Coal Workers' Pneumoconiosis, New Jersey and United States, 2001-2013

| Number and Age-Standardized from Coal Workers' Pneumoconiosis | | | | |
|--|--------|-------|-------|------|
| Year | Number | | Rate* | |
| | NJ | US | NJ | US |
| 2001 | 135 | 6,236 | 19.6 | 28.0 |
| 2002 | 124 | 8,657 | 18.0 | 40.1 |
| 2003 | 131 | 6,590 | 18.3 | 30.3 |
| 2004 | 110 | 5,343 | 15.4 | 24.6 |
| 2005 | 146 | 9,347 | 21.1 | 39.0 |
| 2006 | 134 | 3,853 | 18.2 | 15.4 |
| 2007 | 140 | 2,182 | 18.6 | 8.6 |
| 2008 | 65 | 1,346 | 9.3 | 5.1 |
| 2009 | 47 | 2,224 | 6.3 | 8.9 |
| 2010 | 34 | 670 | 4.6 | 2.6 |
| 2011 | 25 | N/A | 3.3 | N/A |
| 2012 | 29 | N/A | 3.9 | N/A |
| 2013 | 18 | N/A | 2.5 | N/A |

*Rate per million residents N/A: Not Available

NJ Department of Health inpatient hospital discharge data State population estimates from the US Bureau of the Census: www.census.gov/popest/; Year 2000 US Standard Population

Note: Workers' compensation eligibility criteria and availability of data from workers' compensation programs varies among states, prohibiting state-level data from being directly compared to other states or with national estimates.



